

## HA-Tag mouse mAb

www.dia-an.cn

Cat#: 2063

027-59325199


sevice@dia-an.com

Only For Research. Not For Diagnosis.


<b>Synonyms:</b>	YPYDVPDYA
<b>Attribute:</b>	Mouse Monoclonal Antibody
<b>Isotype:</b>	IgG2a
<b>Purity:</b>	Antigen Affinity Purification
<b>Application:</b>	ELISA, WB, IP, IF
<b>Reactivity:</b>	Protein fused with HA tag
<b>Immunogen:</b>	Peptide YPYDVPDYA
<b>Buffer:</b>	PBS with 0.1% sodium azide and 50% glycerol, pH 7.2
<b>Storage:</b>	Store at -20°C. Do not aliquot
<b>Recommended Dilution:</b>	WB: 1:2000-1:10,000 IP: 3ug/sample IF: 1:100-200

### Background:

The HA tag is derived from the HA-molecule corresponding to amino acids 98-106. It has been extensively used as a general epitope tag in expression vectors. Many recombinant proteins have been engineered to express the HA tag, which does not appear to interfere with the bioactivity or the biodistribution of the recombinant protein.



Western Blotting of HA-Tag fused recombinant protein with Anti-HA monoclonal antibody at dilution of 1:2000



Immunofluorescent analysis of 293F cells transfected with the HA-GFP, using 3063R at 1:2000 dilution

We focus on precise protein quantification